

## FIRST RESPONDERS

## OR STAND HOW TO DOFF PPE USING N95 RESPIRATOR


First Responder:		
Trained Observer:	Date:	
<b>PURPOSE:</b> This <b>checklist</b> has been developed to <b>assist First Responders</b> in the <b>proper donning</b> and <b>doffing</b> of Personal Protective Equipment (PPE), and <b>securing</b> on-scene used PPE.		

HOW TO USE: This document should be used by Trained Observers and First Responders in conjunction with:

- The instructional First Responder Ebola Briefing + PPE PowerPoint published November 13, 2014.
- CDC training videos for donning and doffing PPE (www.cdc.gov/vhf/ebola/hcp/ppe-training/).
- CDC Interim Guidance for Emergency Medical Services (EMS) Systems and 9-1-1 Public Safety Answering Points (PSAPs) for Management of Patients with Known or Suspected Ebola Virus Disease in the United States - October 28, 2014. (www.cdc.gov/vhf/ebola/hcp/interim-guidance-emergency-medical-services-systems-911-public-safety-answering-pointsmanagement-patients-known-suspected-united-states.html).
- CDC Guidance on Personal Protective Equipment To Be Used by Healthcare Workers During Management of Patients with Ebola Virus Disease in U.S. Hospitals, Including Procedures for Putting On (Donning) and Removing (Doffing) - October 20, 2014. (www.cdc.gov/vhf/ebola/hcp/procedures-for-ppe.html).

## IMPORTANT NOTES:

- Donning and doffing procedures are different, depending on the type of PPE selected.
- A Trained Observer to verify and document successful compliance with the protocol is required for proper PPE donning and doffing.



Place and double bag used PPE in large (33 gallon or larger) leak-proof plastic bags. Secure at the scene until picked up by a State-provided contractor.		
	Established Work Zone	PPE doffing should be performed in a designated PPE removal area set up <i>prior</i> to donning.  Remember, doffing area should be located as close to the Hot Zone as safely feasible and set up <i>prior</i> to beginning the donning process.
	Prepare to Doff PPE	<ul> <li>□ Before entering PPE removal area, inspect and disinfect any visible contamination on PPE including gloved hands using an EPA-registered disinfectant.</li> <li>□ Step into the pool set up <i>prior</i> to donning and remove any visible contamination using 10% household bleach solution (one [1] part household bleach and nine [9] parts water).</li> <li>□ As used PPE is doffed, drop into leak-proof plastic bags prepared earlier for that purpose.</li> <li>□ Remember: Doffing is your GREATEST RISK of exposure.</li> </ul>
	Step 1. Engage Trained Observer	□ Using a written step-by-step checklist, the doffing process is also conducted under the guidance and supervision of a Trained Observer, who reads aloud each step of the procedure and visually confirms and documents that PPE is removed properly.  □ Trained Observer:  □ Prior to doffing PPE, reminds First Responder to avoid reflexive actions that increase risk of exposure, such as touching the face.  □ Should minimize touching First Responder or his/her PPE during the doffing process.  □ May assist with removal of specific components of PPE, as outlined below.  □ Disinfects outer-gloved hands immediately after handling any First Responder PPE.
	Step 2. Inspect	☐ Inspect PPE for visible contamination, cuts or tears. ☐ Disinfect potentially contaminated PPE using an EPA-registered disinfectant wipe or spray (if facility conditions permit and appropriate regulations are followed).
	Step 3. Disinfect Outer Gloves	Disinfect outer-gloved hands with either an EPA-registered disinfectant wipe OR ABHR. Allow to dry.
	Step 4. Remove Apron (if used)	To avoid contaminating gloves, roll apron from inside (clean side) to outside.
	Step 5. Inspect	☐ Inspect PPE ensemble again. Look for visible contamination, cuts or tears. ☐ IF visibly contaminated, disinfect using EPA-registered disinfectant wipe.
	Step 6. Disinfect Outer Gloves	Use EPA-registered disinfectant wipe or ABHR. Allow to dry.
	Step 7. Remove Boot or Shoe Covers	Sit down to prevent falling. Avoid legs touching each other.  While seated, remove and discard boot or shoe covers.  Manufacturer may have additional instructions for the removal of boot covers.
	Step 8. Disinfect and Remove Outer Gloves	□ Disinfect outer-gloved hands with either an EPA-registered disinfectant wipe OR ABHR. □ Remove and discard outer gloves, taking care NOT to contaminate inner glove. □ Do not snap gloves, causing spray. Peel glove inside out, containing first glove from one hand inside second glove from other hand.

Step 9. Inspect and Disinfect Inner Gloves	□ Inspect the inner gloves' outer surfaces for visible contamination, cuts or tears. □ IF inner glove is visibly soiled, cut or torn □ Disinfect with either an EPA-registered disinfectant wipe □ Perform hand hygiene with ABHR on bare hands. □ Don a clean pair of gloves. □ IF no visible contamination, cuts or tears are identified on inner gloves, disinfect inner-gloved hands with either EPA-registered disinfectant wipe ○R ABHR.
Step 10. Remove Face Shield	Remove full face shield.  — Tilt head slightly forward. — Grab the rear strap and pull it over your head.  Avoid touching the front of the face shield.
Step 11. Disinfect Inner Gloves	Disinfect inner gloves again with either an EPA-registered disinfectant wipe OR ABHR.
Step 12. Remove Hood	Unfasten hood. Gently remove and discard.  Trained Observer may assist with unfastening hood.
Step 13. Disinfect Inner Gloves	Disinfect inner gloves again with either an EPA-registered disinfectant wipe OR ABHR.
Step 14. Remove Coverall	☐ Tilt head back. CAREFULLY locate zipper or fasteners. ☐ Using two (2) fingers of one (1) hand only, unzip or unfasten coverall completely. ☐ Roll coverall downward, turning inside (clean side) out. ☐ Avoid contact of scrubs or disposable garments with outer surface of coverall. ☐ Touch ONLY the inside (clean side) of the coverall.
Step 15. Disinfect and Change Inner Gloves	Disinfect inner-gloved hands with either an EPA-registered disinfectant wipe OR ABHR.  — CAREFULLY remove and discard inner gloves.   — Do NOT contaminate bare hands.  — Perform hand hygiene with ABHR.  — Don NEW pair of gloves.
Step 16. Remove N95 Respirator	☐ Tilt head forward. Grasp the bottom tie or elastic strap, then the top tie or elastic strap. ☐ Remove — without touching the front of the N95 respirator. ☐ Discard N95 respirator.
Step 17. Disinfect Inner Gloves	Disinfect inner gloves again with either an EPA-registered disinfectant wipe OR ABHR.
Step 18. Disinfect Washable Shoes	Sit on a new, clean surface (e.g., second clean chair, clean side of a bench).  Using an EPA-registered disinfectant wipe, wipe down every external surface, including soles, of washable shoes. Avoid touching ankles.
Step 19. Disinfect and Remove Inner Gloves	Use EPA-registered disinfectant wipe OR ABHR.  CAREFULLY remove and discard inner gloves. Do not snap gloves, causing spray. Peel glove inside out, containing first glove from one hand inside second glove from other hand.  Do NOT contaminate bare hands. Do not touch face or bare skin.
Step 20. Perform Hand Hygiene	Perform hand hygiene with ABHR.
Step 21. Inspect	Perform a <b>final inspection</b> for any indication of <b>contamination</b> of <b>scrubs</b> or <b>disposable garments</b> .  IF contamination is identified, <b>immediately</b> notify supervisor <b>before exiting</b> PPE removal area.
Step 22. Scrubs	First Responder can leave PPE removal area wearing dedicated washable footwear and surgical scrubs or disposable garments.
Step 23. Shower	Showers are recommended for those  — Performing high-risk patient care (e.g., exposed to large quantities of blood, body fluids or excreta).  — Spending extended periods of time with an Ebola patient.
Step 24. Protocol Evaluation/ Medical Assessment	☐ Either the infection preventionist or occupational safety and health coordinator or their designee should meet with the First Responder to review patient care activities performed to identify any concerns about care protocols and to record First Responder's level of fatigue.

N95 Respirator Doffing Page 2 of 2









